

**DE NOVO OF FEMORAL CONDYLE  
REHABILITATION PROTOCOL**

	<b>WEIGHTBEARING</b>	<b>BRACE</b>	<b>ROM</b>	<b>EXERCISES</b>
<b>PHASE I</b> 0-6 weeks	Non-WB	<b>0-2 week:</b> Locked in Full extension at all times  Off for CPM and exercise only  Discontinue after 2 wks	<b>0-6 weeks:</b> Use CPM for 6 hours/day, beginning at 0- 40°; advance 5-10° daily as tolerated	<b>0-2 weeks:</b> Quad sets, SLR, calf pumps, passive leg hangs to 90° at home  <b>2-6 weeks:</b> PROM/AAROM to tolerance, patella and tibiofibular joint mobs, quad, hamstring, and glute sets, SLR, side-lying hip and core
<b>PHASE II</b> 6-8 weeks	Advance 25% weekly until full	None	Full	Advance Phase I exercises
<b>PHASE III</b> 8-12 weeks	Full	None	Full	Gait training, begin closed chain activities: wall sits, shuttle, mini-squats, toe raises  Begin unilateral stance activities, balance training
<b>PHASE IV</b> 12 weeks - 6 months	Full	None	Full	Advance Phase III exercises; maximize core/glutes, pelvic stability work, eccentric hamstrings  May advance to elliptical, bike, pool as tolerated
<b>PHASE V</b> 6-12 months	Full	None	Full	Advance functional activity  Return to sport-specific activity and impact when cleared by MD after 8



P: 405.885.8195 F: 405.218.2535 E: [ortho@jtglomsetmd.com](mailto:ortho@jtglomsetmd.com)  
[www.jtglomsetmd.com](http://www.jtglomsetmd.com)

				mos
--	--	--	--	-----