

P: 405.885.8195 F: 405.218.2535 E: ortho@jtglomsetmd.com www.jtglomsetmd.com

## POSTEROLATERAL CORNER RECONSTRUCTION REHABILITATION PROTOCOL

	WEIGHTBEARING	BRACE	ROM	EXERCISES
DUACE :				
PHASE I 0-6 weeks	Heel touch WB in brace *	O-2 weeks: Locked in full extension for ambulation and sleeping  2-6 weeks:	0-2 weeks: 0- 45 2-6 weeks: Advance slowly 0-90	Quad sets, patellar mobs, gastroc/soleus stretch  SLR w/ brace in full extension until quad strength prevents extension lag
		Unlocked for ambulation 0-90, remove for sleeping**		Side-lying hip/core  Hamstrings avoidance until 6 wks post-op
PHASE II 6-12 weeks	Advance 25% weekly until full by 8 wks	Discontinue at 6 wks if no extension lag	Full	Begin toe raises, closed chain quads, balance exercises, hamstring curls, stationary bike, step-ups, front and side planks; advance hip/core
PHASE III 12-16 weeks	Full	None	Full	Advance closed chain strengthening  Progress proprioception activities  Begin stairmaster, elliptical and running straight ahead at 12 weeks
PHASE IV 16-24 weeks	Full	None	Full	16 wks: Begin jumping  20 wks: Advance to sprinting, backward running, cutting/pivoting/changing direction, initiate plyometric program and sport-specific drills
PHASE V > 6 months	Full	None	Full and pain free	Gradual return to sports participation after completion of FSA***  Maintenance program based on FSA



P: 405.885.8195 F: 405.218.2535 E: ortho@jtglomsetmd.com www.jtglomsetmd.com

- \*Modified with concomitantly performed meniscus repair/transplantation or articular cartilage procedure
- \*\*Brace may be removed for sleeping after first post-operative visit (day 7-10)
- \*\*\*Completion of FSA (Functional Sports Assessment) not mandatory, but recommended at 22-24 wks post-op for competitive athletes returning to play after rehab